

$O(d+1, d+1)$ enhanced double field theory

Olaf Hohm,^a Edvard T. Musaev^{b,c} and Henning Samtleben^d

^a*Simons Center for Geometry and Physics, Stony Brook University,
Stony Brook, NY 11794-3636, U.S.A.*

^b*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),
Am Mühlenberg 1, DE-14476 Potsdam, Germany*

^c*General Relativity Department, Institute of Physics, Kazan Federal University,
Kremlevskaya 16a, 420111, Kazan, Russia*

^d*Univ Lyon, ENS de Lyon, Univ Claude Bernard Lyon 1, CNRS, Laboratoire de Physique,
F-69342 Lyon, France*

E-mail: ohohm@scgp.stonybrook.edu, edvard.musaev@aei.mpg.de,
henning.samtleben@ens-lyon.fr

ABSTRACT: Double field theory yields a formulation of the low-energy effective action of bosonic string theory and half-maximal supergravities that is covariant under the T-duality group $O(d, d)$ emerging on a torus T^d . Upon reduction to three spacetime dimensions and dualisation of vector fields into scalars, the symmetry group is enhanced to $O(d+1, d+1)$. We construct an enhanced double field theory with internal coordinates in the adjoint representation of $O(d+1, d+1)$. Its section constraints admit two inequivalent solutions, encoding in particular the embedding of $D=6$ chiral and non-chiral theories, respectively. As an application we define consistent generalized Scherk-Schwarz reductions using a novel notion of generalized parallelization. This allows us to prove the consistency of the truncations of $D=6$, $\mathcal{N}=(1, 1)$ and $D=6$, $\mathcal{N}=(2, 0)$ supergravity on $\text{AdS}_3 \times \mathbb{S}^3$.

KEYWORDS: Bosonic Strings, M-Theory, String Duality, String Field Theory

ARXIV EPRINT: [1707.06693](https://arxiv.org/abs/1707.06693)